

Particle dependence of elliptic flow in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV

Paul Sorensen[†] for the STAR collaboration[‡]

[†] University of California, Los Angeles, California 90095

E-mail: sorensen@physics.ucla.edu

Abstract. The elliptic flow parameter (v_2) for K_S^0 and $\Lambda + \bar{\Lambda}$ has been measured at mid-rapidity in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV by the STAR collaboration. The v_2 values for both K_S^0 and $\Lambda + \bar{\Lambda}$ saturate at moderate p_T , deviating from the hydrodynamic behavior observed in the lower p_T region. The saturated v_2 values and the p_T scales where the deviation begins are particle dependent. The particle-type dependence of v_2 shows features expected from the hadronization of a partonic ellipsoid by coalescence of co-moving quarks. These results will be discussed in relation to the nuclear modification factor (R_{CP}) which has also been measured for K_S^0 and $\Lambda + \bar{\Lambda}$ by the STAR collaboration.

Submitted to: *J. Phys. G: Nucl. Phys.*

PACS numbers: 25.75.-q, 25.75.Ld, 25.75.Dw

1. Introduction

The elliptic component of the event-wise azimuthal anisotropy of particle production (i.e. *elliptic flow* or v_2) is thought to probe the early stages of relativistic heavy ion collisions [2]. Measurements of v_2 for identified particles [3, 4] and charged hadrons [5, 6] at the Relativistic Heavy Ion Collider (RHIC) indicate a conversion of spatial anisotropy to momentum anisotropy near the hydrodynamical limit [2, 7, 8]. For p_T greater than 2 GeV/c, however, the charged hadron v_2 deviates from hydrodynamical calculations and saturates at a large value approximately independent of p_T up to 6 GeV/c [6]. The measurements of v_2 for K_S^0 and $\Lambda + \bar{\Lambda}$ also indicate a saturation at moderate p_T . Models using large parton energy loss [9, 10] and transport opacity [11] have been discussed in relation to the saturation and centrality dependence of charged hadron v_2 at large p_T . A saturated v_2 could also arise if the extent of the p_T region where *soft processes* dominate the spectrum is particle dependent [12]. In this case the hyperon v_2 may continue to rise at intermediate p_T (perhaps following hydrodynamic model calculations) while v_2 of the less massive meson decreases. It has also been suggested that if a partonic state

[‡] For a complete collaboration list see [1]

exists prior to hadronization, the process of particle formation at moderately high p_T , by string fragmentation, parton fragmentation [13] or quark coalescence [14, 15, 16, 17], may lead to a dependence of v_2 and R_{AA} on particle type. As such, it's possible that these measurements will provide information on the existence and nature of an early partonic state.

In this Letter, we report the measurement of v_2 at mid-rapidity, $|y| \leq 1.0$, for K_S^0 and lambda (Λ) + antilambda ($\bar{\Lambda}$) with $0.4 < p_T < 6.0$ GeV/c and $0.6 < p_T < 6.0$ GeV/c respectively in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

2. Analysis and Results

This analysis uses 1.6×10^6 minimum-bias trigger events and 1.5×10^6 central trigger events detected in the STAR detector system [18]. The particles, K_S^0 , Λ and $\bar{\Lambda}$, were identified from the charged daughter tracks produced in the decays $K_S^0 \rightarrow \pi^+ + \pi^-$, $\Lambda \rightarrow p + \pi^-$ and $\bar{\Lambda} \rightarrow \bar{p} + \pi^+$. A detailed description of the analysis, such as track finding, decay vertex topology cuts, and the estimation of detection efficiency, is given in Refs. [4, 19].

We use the yield as a function of $(\phi_{ij} - \Psi_j^R)$ to calculate $v_2 = \langle \cos[2(\phi_{ij} - \Psi_j^R)] \rangle$, where ϕ_{ij} is the azimuthal emission angle of particle i in event j and Ψ_j^R is the reaction plane angle for event j , where, to remove autocorrelations, the decay daughter tracks associated with particle i are excluded from its calculation. Within statistical errors, the Λ v_2 is the same as $\bar{\Lambda}$ v_2 , so they are summed together.

Table 1. The systematic errors for minimum bias v_2 from non-flow effects (n-f) and background contamination (bkg). The values represent the *absolute* errors. The p_T resolution, $\delta p_T/p_T$ is also listed.

	K_S^0			$\Lambda + \bar{\Lambda}$		
p_T (GeV/c)	1.0	2.5	4.0	1.0	2.5	4.0
v_2 (bkg)	+0.000 -0.001	+0.001 -0.007	+0.003 -0.018	+0.001 -0.007	+0.005 -0.001	+0.005 -0.001
v_2 (n-f)	+0.00 -0.01	+0.00 -0.04	+0.00 -0.03	+0.00 -0.01	+0.00 -0.04	+0.00 -0.04
$\delta p_T/p_T$	0.016	0.027	0.037	0.016	0.027	0.037

Possible sources of systematic error in the calculation of v_2 are correlations unrelated to the reaction plane (non-flow effects), uncertainties in the extraction of yields from the invariant mass distributions, the particle momentum resolution ($\delta p_T/p_T$), and biases introduced by the cuts used in the analysis. Table 1 lists the dominant systematic errors for three transverse momenta. The non-flow systematic error is dominant. The non-flow effects for charged particle v_2 are discussed in Refs. [6, 5] but, the particle dependence of these effects has not been measured. We assume a similar magnitude of non-flow contribution to $\Lambda + \bar{\Lambda}$ and K_S^0 v_2 . A 4-particle cumulant analysis of $\Lambda + \bar{\Lambda}$ and K_S^0 v_2 will be less sensitive to non-flow effects but, to be conclusive, will require a larger data sample than is currently available.




Figure 1. The elliptic flow of K_S^0 and $\Lambda + \bar{\Lambda}$ as a function of p_T for 30–70% (a), 5–30% (b), 0–5% (c) and 0–80% (d) of the collision cross section. The error bars shown are statistical errors only.

Fig. 1 shows v_2 of K_S^0 and $\Lambda + \bar{\Lambda}$ as a function of p_T for the centrality intervals, 30–70% (a), 5–30% (b), 0–5% (c), and 0–80% (d) of the geometrical cross section. The p_T dependence of the v_2 for all the centrality bins has a similar trend. There is a saturation and particle dependence at moderate p_T for each of the centrality intervals and, as such, the saturation for the minimum-bias v_2 (0–80%) cannot be due to the superposition of drastically different p_T dependencies in various centrality bins. This measurement establishes the saturation and particle dependence of v_2 at the moderate to high p_T region.




Figure 2. (Top panel) The elliptic flow of K_S^0 and $\Lambda + \bar{\Lambda}$ as a function of $m_T - m_0$ for 30–70% (a), 5–30% (b) and 0–5% (c) of the collision cross section. The error bars shown are statistical errors only. (bottom panel) The same scaled by the initial eccentricity.

Fig. 2 and Fig. 3 show the same data as Fig. 1 but plotted versus $m_T - m_0$ which is




Figure 3. The elliptic flow of K_S^0 and $\Lambda + \bar{\Lambda}$ as a function of p_T (top panel) and $m_T - m_0$ (bottom panel) for 0–80% of the collision cross section. The error bars shown are statistical errors only.

approximately the kinetic energy of the hadron. In the low momentum region the K_S^0 and $\Lambda + \bar{\Lambda}$ v_2 appears to fall on a single straight line. The hydrodynamic calculations in Fig. 3 seem to capture this trend. In the bottom panels of Fig. 2 we’ve scaled the v_2 values by the eccentricity of the overlap region for the various centralities. Hydrodynamic models predict that v_2 should scale with this initial eccentricity.

In Fig. 4 (top) we show $v_2(p_T)$ for K_S^0 , $\Lambda + \bar{\Lambda}$, and charged hadrons [20] along with hydrodynamic model calculations of v_2 for identified particles [7]. Below $p_T \sim 1.2$ GeV/c v_2 is consistent with the calculations and in agreement with the previous results for K_S^0 and $\Lambda + \bar{\Lambda}$ at $\sqrt{s_{NN}} = 130$ GeV [4]. Contrary, however, to hydrodynamical calculations, where at a given p_T heavier particles will have smaller v_2 values, the measured v_2 of the heavier hyperon saturates at a value significantly *larger* than the v_2 of the lighter K_S^0 meson. The p_T scale where the measured v_2 deviates from the hydrodynamical prediction is particle dependent with the hyperon v_2 following the prediction up to $p_T \sim 2.0$ GeV/c while the K_S^0 v_2 deviates much sooner.

The ratio (R_{CP}) of the yields in central and peripheral collisions scaled by the number of binary nucleon-nucleon collisions (N_{binary}), may also be sensitive to the effects of energy loss and hadronization via parton coalescence [16, 9]. In Fig. 4 (bottom) we show R_{CP} for K_S^0 , $\Lambda + \bar{\Lambda}$ and charged hadrons using the centrality intervals 0–5% (central) and 60–80% (peripheral) [21, 22]. The charged hadron spectrum at $p_T > 2$ GeV/c for the 60–80% centrality bin approximately follows binary collision scaling without medium modification [22]. As such, when this bin is used, R_{CP} approximates R_{AA} . The bands in Fig. 4 represent the expected values of R_{CP} for N_{binary} and N_{part} scaling including systematic variations within the calculation [22].

For $p_T < 5$ GeV/c, the K_S^0 and $\Lambda + \bar{\Lambda}$ yields are suppressed (relative to N_{binary} scaling) by different magnitudes. In addition, the p_T scale associated with the onset suppression has a dependence on particle-type that is similar to the dependence in v_2 for the onset of saturation. At $p_T \sim 5.0$ GeV/c, R_{CP} values for K_S^0 and $\Lambda + \bar{\Lambda}$ are both approaching the value of the charged hadron R_{CP} .




Figure 4. Minimum-bias v_2 (top panel) and R_{CP} (bottom panel) for K_S^0 and $\Lambda + \bar{\Lambda}$. The v_2 errors are statistical only. The widths of the gray bands represent uncertainties in the model calculation of N_{binary} and N_{part} . Charged hadron v_2 and R_{CP} are also shown.

3. Discussion

Although R_{CP} depends only on the yield in the central and peripheral bins, and the v_2 in Fig. 4 is from a minimum-bias centrality interval, the two parameters may be intimately related. The differential elliptic flow ($v_2(p_T)$) measures the ratio of the p_T spectrum of particles emitted in the direction of the reaction plane (in-plane) to that of particles emitted perpendicular the reaction plane (out-of-plane). In hydrodynamic models, we expect the pressure gradient to be larger in the in-plane direction than the out-of-plane direction. In this case, v_2 will be the ratio of a p_T spectrum with more hydrodynamic flow to one with less flow. As such, taking the ratio (R_{CP}) of a p_T spectrum that exhibits large flow (central) and one exhibiting less flow (peripheral) *should* lead to a very similar p_T and particle-type dependence. A detailed study of v_2 and R_{CP} for identified particles should reveal the extent (in p_T and centrality) to which hydrodynamical models are valid.

A *surface emission* scenario, where partons traversing a dense medium experience large energy losses, has been discussed in relation to the large, p_T independent v_2 measured for charged hadrons [10]. This mechanism should also lead to a suppression of particle production in central Au+Au collisions. This scenario, however, is inconsistent with STAR measurements of v_2 and R_{CP} for K_S^0 and $\Lambda + \bar{\Lambda}$. The smaller suppression manifested in the $\Lambda + \bar{\Lambda}$ R_{CP} contradicts the larger azimuthal anisotropies manifested

in $\Lambda + \bar{\Lambda}$ v_2 values. In addition, calculations based on a surface emission model [10] cannot produce v_2 values as large as those measured.

The absence of a net suppression of $\Lambda + \bar{\Lambda}$ for p_T from 1.8–3.5 GeV/c in central Au+Au collisions could also indicate the presence of dynamics beyond the framework of parton energy loss followed by fragmentation. The stronger dependence on centrality (and thus parton density) for baryon production indicated by the larger R_{CP} would naturally be expected from multi-parton mechanisms such as gluon junctions [23], quark coalescence [14], or recombination [16]. Within the framework of these models, the measured v_2 and R_{CP} features may reflect the anisotropy and hadronization properties of the bulk quark matter. Fig. 5 shows v_2 of K_S^0 and $\Lambda + \bar{\Lambda}$ as a function of p_T where the v_2 and p_T values have been scaled by the number of constituent quarks (n). Above $p_T/n \sim 0.8$ GeV/c, the v_2/n vs p_T/n is the same, within errors, for both species. In a scenario where hadrons at intermediate p_T ($\sim 1 - 5$ GeV/c) are predominantly formed from bulk partonic matter by quark coalescence, e.g. Ref. [14], v_2/n should reveal the v_2 developed by partons prior to the hadronic phase. The verification of this scenario, to the exclusion of other possible explanations, would be strong evidence for the formation of a quark-gluon plasma at RHIC.




Figure 5. The v_2 parameter for K_S^0 and $\Lambda + \bar{\Lambda}$ scaled by the number of constituent quarks (n) and plotted versus p_T/n .

The scenario discussed in Ref. [12], where soft processes dominate the hyperon v_2 up to a higher p_T than mesons, could also lead to a particle dependence for R_{CP} that is qualitatively consistent with these measurements. Quantitative calculations of v_2 and predictions for R_{CP} from this scenario, however, are still needed. Up-coming measurements of R_{CP} for identified particles in d + Au collisions at RHIC will also make it possible to study the effect of initial state interactions on R_{CP} .

4. Summary

We have reported the measurement of v_2 for p_T up to ~ 6.0 GeV/c for K_S^0 and $\Lambda + \bar{\Lambda}$ from Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. For $p_T < 1.2$ GeV/c, hydrodynamic model calculations agree well with the p_T and mass dependence of the measured v_2 . At this low momentum region K_S^0 and $\Lambda + \bar{\Lambda}$ v_2 lie on a single straight line when plotted versus $m_T - m_0$. In the moderate p_T region, however, the particle type and p_T dependence of v_2 suggests hydrodynamics no longer describes the collision dynamics. The value of v_2 for K_S^0 saturates earlier and at a lower value than the $\Lambda + \bar{\Lambda}$ v_2 . Measurements of R_{CP} show that the suppression of particle production in central collisions depends on particle-type in a similar way. The measurement of the particle-type and p_T dependence of v_2 and R_{CP} at moderate p_T may provide a unique means to establish the existence (and study the properties) of a quark-gluon plasma that may be formed in collisions at RHIC.

Acknowledgments: This work has been partially supported by NSF grant PHY-03-11859.

5. References

- [1] H. Caines (for the STAR Collaboration), these proceedings.
- [2] J.-Y. Ollitrault, Phys. Rev. D **46**, 229(1992); H. Sorge, Phys. Rev. Lett. **82**, 2048(1999).
- [3] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **87**, 182301 (2001); S. Esumi (for the PHENIX Collaboration), e-print:nucl-ex/0210012.
- [4] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **89**, 132301 (2002); P. Sorensen (for the STAR Collaboration), J. Phys. G **28**, 2089 (2002).
- [5] C. Adler *et al.* (STAR Collaboration), Phys. Rev. C **66**, 034904 (2002).
- [6] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **90**, 032301 (2003).
- [7] P. Huovinen, P.F. Kolb, U. Heinz, P.V. Ruuskanen, and S.A. Voloshin, Phys. Lett. B **503**, 58(2001).
- [8] D. Teaney, J. Lauret, and E.V. Shuryak, Phys. Rev. Lett. **86**, 4783(2001).
- [9] E. Wang and X. N. Wang, Phys. Rev. Lett. **89**, 162301 (2002);
- [10] E. V. Shuryak, Phys. Rev. C **66**, 027902 (2002).
- [11] D. Molnar and M. Gyulassy, Nucl. Phys. A **697**, 495 (2002) [Erratum-ibid. A **703**, 893 (2002)]
- [12] M. Gyulassy, I. Vitev, X. N. Wang and P. Huovinen, Phys. Lett. B **526**, 301 (2002).
- [13] Z.W. Lin and C.M. Ko, Phys. Rev. C **65**, 034904(2002).
- [14] D. Molnar and S. A. Voloshin, e-print:nucl-th/0302014.
- [15] Z. W. Lin and C. M. Ko, Phys. Rev. Lett. **89**, 202302 (2002)
- [16] R. J. Fries, B. Müller, C. Nonaka and S. A. Bass, e-print:nucl-th/0301087.
- [17] V. Greco, C. M. Ko and P. Levai, e-print:nucl-th/0301093.
- [18] C. Adler *et al.* (STAR Collaboration), Nucl. Instr. Meth. A **499**, 624 (2003).
- [19] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **89**, 092301 (2002); K. Safarik, I. Kraus, J. Newby and P. Sorensen, AIP Conf. Proc. **631**, 377 (2003).
- [20] K. Filimonov (for the STAR Collaboration), e-print:nucl-ex/0210027.
- [21] H. Long (for the STAR Collaboration), these proceedings.
- [22] J. L. Klay (for the STAR Collaboration), e-print:nucl-ex/0210026.
- [23] S.E. Vance and M. Gyulassy, Phys. Rev. Lett. **83**, 1735 (1999).